

Unitarity Triangle *Without* Semileptonic Decays

E. Lunghi¹ and Amarjit Soni²

¹*Physics Department, Indiana University, Bloomington, IN 47405*

²*Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

Use of semi-leptonic decays has become standard in constraining the Unitarity Triangle. Bearing in mind that precise calculations of these are very challenging, we propose an entirely new approach. In particular the $|V_{cb}| + \varepsilon_K$ constraint, which depends extremely sensitively on $|V_{cb}|$ in the traditional method, is replaced by the interplay between ε_K , $\text{BR}(B \rightarrow \tau\nu)$ and ΔM_{B_s} . It is found that even in this method tensions with the Standard Model persist at the $\sim 1.8\sigma$ level. Furthermore, improvements on the $B \rightarrow \tau\nu$ branching ratio and on the lattice determination of $f_{B_s} \hat{B}_s^{1/2}$ can increase the effectiveness of this method significantly.

PACS numbers: 12.15.Hh, 11.30.Hv

The next big step in our understanding of particle physics will be the uncovering of the electro-weak symmetry breaking (EWSB) mechanism. The present and upcoming collider experiments (Fermilab and LHC) will be able to test the Standard Model (SM) Higgs mechanism. New physics is widely expected at around the TeV scale if the Higgs mass is not to receive large radiative corrections and require severe fine-tuning. A stringent constraint on the SM mechanism of EWSB is the tight structure of flavor changing (FC) interactions: tree-level FC neutral currents are forbidden and charged currents are controlled by the Cabibbo–Kobayashi–Maskawa (CKM) [1] mixing matrix

$$V = \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix}.$$

Within the SM, the CKM matrix is the only source of FC interactions and of CP violation. There is no reason, in general, to expect that new physics (needed to stabilize the Higgs mass) at the TeV scale will be in the basis wherein the quark mass matrix is diagonal. This reasoning gives rise to another fundamental problem in particle physics, namely the flavor puzzle i.e. unless the scale of new physics is larger than 10^3 TeV it causes large FCNC especially for the $K - \bar{K}$ system. Thus flavor physics provides constraints on models of new physics upto scales that are much much larger than what is accessible to direct searches at colliders such as the Tevatron or the LHC. Flavor physics is therefore expected to continue to provide crucial information for the interpretation of any physics that LHC may find.

In this decade significant progress has been made in our understanding of flavor physics, thanks in large part to the spectacular performance of the two asymmetric B-factories. For the first time, we learned that the CKM-paradigm of CP violation is able to simultaneously account for the observed CP-violation in the K and B-systems up to an accuracy of O(20%). Impressive as this is, it is still important to understand that this leaves

a lot of room for new physics. Indeed, as more data from B-factories became available and also as the accuracy in some key theoretical calculations was attained, several 2-3 σ hints of new physics have emerged. While this clearly does not represent an unambiguous signal for new physics, it does mean that efforts need to continue both on the experiment and on the theory front to seek greater clarity with regard to these anomalies.

In this context use of semileptonic decays in all traditional analysis of the Unitarity Triangle (UT) to date is a concern. The inclusive $b \rightarrow u$ transitions are not governed by any symmetry and as a result are a special challenge for continuum methods. Exclusive decays are in principle amenable to the lattice and steady, but unfortunately rather slow, progress is being made. The fact that for both $b \rightarrow c$ and for $b \rightarrow u$ inclusive and exclusive methods disagree by $\approx 2\sigma$ casts a shadow of doubt on the results of the UT analysis. This is especially aggravated by the fact that use of the input from ε_K , representing the indirect CP violation from the $K_L \rightarrow \pi\pi$, into the UT is exceedingly sensitive to V_{cb} , scaling as the fourth power. These observations motivate us to seek alternate approaches, which we will provide herein, though we want to emphasize that we are not suggesting that efforts to improve our understanding of semi-leptonic decays be discontinued but rather that alternate methods of analysis for the UT could be very valuable and should also be developed.

Recent improvements especially in the lattice calculation of B_K [2, 3, 4, 5], led to the appearance of a $\sim 2\sigma$ tension that can be interpreted as new physics in B_d and/or in K mixing [6, 7, 8, 9, 10]. The interplay of ε_K with $\Delta M_{B_s}/\Delta M_{B_d}$ and $S_{\psi K}$ (time-dependent CP asymmetry in $B \rightarrow J/\psi K_s$) is at the heart of the tension. The inclusion of $|V_{ub}|$ from semileptonic $b \rightarrow u\ell\nu$ decays, tends to favor a scenario with new physics in kaon mixing [11]. An important difficulty with these analyses is the long standing discrepancy between the extraction of $|V_{cb}|$ and $|V_{ub}|$ from exclusive and inclusive semileptonic decays alluded to above. From the inspection of Table I, one sees that inclusive and exclusive determinations dif-

$ V_{cb} _{\text{excl}} = (38.6 \pm 1.2)10^{-3}$	$\eta_1 = 1.51 \pm 0.24$
$ V_{cb} _{\text{incl}} = (41.31 \pm 0.76)10^{-3}$	$\eta_2 = 0.5765 \pm 0.0065$
$ V_{cb} _{\text{incl+excl}} = (40.3 \pm 1.0)10^{-3}$	$\eta_3 = 0.47 \pm 0.04$
$ V_{ub} _{\text{excl}} = (34.2 \pm 3.7)10^{-4}$	$\eta_B = 0.551 \pm 0.007$
$ V_{ub} _{\text{incl}} = (40.1 \pm 2.7 \pm 4.0)10^{-4}$	$\xi = 1.23 \pm 0.04$
$ V_{ub} _{\text{incl+excl}} = (36.4 \pm 3.0)10^{-4}$	$\lambda = 0.2255 \pm 0.0007$
$\Delta M_{B_d} = (0.507 \pm 0.005) \text{ ps}^{-1}$	$\alpha = (89.5 \pm 4.3)^\circ$
$\Delta M_{B_s} = (17.77 \pm 0.12) \text{ ps}^{-1}$	$S_{\psi K_S} = 0.672 \pm 0.024$
$\varepsilon_K = (2.229 \pm 0.012) \times 10^{-3}$	$\gamma = (78 \pm 12)^\circ$
$m_c(m_c) = (1.268 \pm 0.009) \text{ GeV}$	$\widehat{B}_K = 0.725 \pm 0.027$
$m_{t,\text{pole}} = (172.4 \pm 1.2) \text{ GeV}$	$f_B = (192.8 \pm 9.9) \text{ MeV}$
$f_{B_s} \sqrt{\widehat{B}_s} = (275 \pm 19) \text{ MeV}$	$f_K = (155.8 \pm 1.7) \text{ MeV}$
$\mathcal{B}_{B \rightarrow \tau \nu} = (1.43 \pm 0.37)10^{-4} [16]$	$\kappa_\varepsilon = 0.92 \pm 0.01$

TABLE I: Inputs used in the fit. References to the original experimental and theoretical papers and the description of the averaging procedure can be found in Ref. [11]. Statistical and systematic errors are combined in quadrature. We adopt the averages of Ref. [11] for all quantities with the exception of $|V_{ub}|$, ξ and $f_{B_s} \sqrt{\widehat{B}_s}$ (see text).

fer at the 2σ level. While Ref. [7] demonstrated that $|V_{ub}|$ can be dropped from the fit without affecting the observed tension, it is usually believed that $|V_{cb}| \simeq A\lambda^2$ from semileptonic decays is essential in order to use ε_K (because of its A^4 dependence).

Bearing all this in mind, in this letter we propose a new approach to the UT analysis, wherein no use of semileptonic decays is made. In particular, we show that the traditional use of $\varepsilon_K + |V_{cb}|$ combination can be effectively replaced by the interplay between ε_K , ΔM_{B_s} and $\text{BR}(B \rightarrow \tau \nu)$. We find that even after removing all input to UT analysis from semileptonic decays, the 2σ tension in the UT fit with the SM survives. More importantly, every experimental and theoretical input to this analysis is now theoretically clean and under very good control. The latter point is quite important, because many of the hints for new physics that come from precision studies tend to have some problems. For instance, the muon anomalous magnetic moment ($(g-2)_\mu$) tension relies on the non-perturbative estimation of light-by-light scattering contributions, the use of $e^+e^- \rightarrow \text{hadrons}$ to calculate the hadronic vacuum polarization tensor of the photon and on isospin corrections to τ decays data. Another example is the $(2-3)\sigma$ tensions in the direct CP asymmetries in $B \rightarrow K\pi$ as well as in time-dependent CP asymmetries $B \rightarrow \psi K$ versus the penguin modes such as $B \rightarrow \phi K$ or $\eta' K$ etc. These rely on model dependent QCD treatment of power corrections and non-factorizable effects which are very difficult to ascertain reliably. A very important exception is the 2.2σ evidence for a CP violating phase in B_s mixing, whose non-zero value would be an extremely clean evidence for new physics [12, 13, 14, 15].




FIG. 1: Standard unitarity triangle fit. The contour is obtained using V_{cb} , V_{ub} , ε_K , $B \rightarrow \tau \nu$, γ , ΔM_{B_s} and ΔM_{B_d} .

Present status of the UT fit. The standard technique to extract the parameters A , ρ and η relies on a simultaneous fit of the following observables: the mixing induced CP violation in the kaon system (ε_K), the mass differences in the B_d and B_s systems ($\Delta M_{B_{d,s}}$), the CP asymmetries in $B \rightarrow J/\psi K_S$ ($S_{\psi K}$), $B \rightarrow (\pi\pi, \rho\rho, \rho\pi)$ (α) and $B \rightarrow D^{(*)}K^{(*)}$ (γ) modes, the rates of semileptonic $b \rightarrow (c, u)\ell\nu$ decays ($|V_{cb}|$ and $|V_{ub}|$) [17] and the $B \rightarrow \tau\nu$ branching ratio. A complete description of these observables and of the possible statistical techniques that could be employed can be found in Refs. [18, 19]. We follow the approach of Refs. [7, 10] and utilize the averages calculated in Ref. [11] with some exceptions: we include inclusive $|V_{ub}|$ albeit with an additional 10% model uncertainties [20]; we take a simple (not weighted) average of the determinations of ξ from Fermilab/MILC [21] and HPQCD [22]. Also we take the central value of $f_{B_s} \sqrt{\widehat{B}_s}$ from Ref. [11] but adopt the uncertainty quoted in Ref. [22]. We adopt this conservative stance to show that the impact of the approach we champion in this letter remains largely unaffected even if the lattice errors are not as small as currently claimed in the literature.

We summarize the inputs we use in Table I. Below we first present explicitly only those formulas that are relevant to the traditional analysis which uses semi-leptonic decays:

$$\Delta M_{B_s} = \chi_s f_{B_s}^2 \widehat{B}_{B_s} A^2 \lambda^4 \quad (1)$$

$$\Delta M_{B_d} = \chi_d f_{B_d}^2 \widehat{B}_{B_d} A^2 \lambda^6 (\eta^2 + (-1 + \rho)^2) \quad (2)$$

$$\frac{\Delta M_{B_s}}{\Delta M_{B_d}} = \frac{m_{B_s}}{m_{B_d}} \frac{\xi^2 \lambda^{-2}}{\eta^2 + (-1 + \rho)^2} \quad (3)$$

$$|\varepsilon_K| = 2\chi_\varepsilon \widehat{B}_K \kappa_\varepsilon \eta \lambda^6 \left(A^4 \lambda^4 (\rho - 1) \eta_2 S_0(x_t) + A^2 (\eta_3 S_0(x_c, x_t) - \eta_1 S_0(x_c)) \right) \quad (4)$$

$$\text{BR}(B \rightarrow \tau \nu) = \chi_\tau f_B^2 |V_{ub}|^2 \simeq \chi_\tau f_B^2 A^2 \lambda^6 (\rho^2 + \eta^2) \quad (5)$$

where we expanded in λ and defined

$$\chi_q = G_F^2 m_W^2 m_{B_q} \eta_B S_0(x_t)/(6\pi^2), \quad (6)$$

$$\chi_\varepsilon = (G_F^2 m_W^2 f_K^2 m_K)/(12\sqrt{2}\pi^2 \Delta m_K^{\text{exp}}), \quad (7)$$

$$\chi_\tau = G_F^2 m_\tau^2 m_{B^+}/(8\pi\Gamma_{B^+})(1 - m_\tau^2/m_{B^+}^2)^2. \quad (8)$$

The 68% C.L. allowed regions in the (ρ, η) plane are shown in Fig. 1, where we show explicitly that the ε_K , $B \rightarrow \tau\nu$ (pink) and $|V_{ub}|$ (yellow) constraints require $|V_{cb}|$ in order to be drawn independently. In particular we obtain:

$$|V_{ub}|_{\text{fit}} = (3.61 \pm 0.13) \times 10^{-3}, \quad (9)$$

$$\text{BR}(B \rightarrow \tau\nu)_{\text{fit}} = (0.87 \pm 0.11) \times 10^{-4}, \quad (10)$$

$$[\sin 2\beta]_{\text{fit}} = 0.766 \pm 0.036, \quad (11)$$

where each quantity is extracted by removing the corresponding direct determination from the fit (for V_{ub} this means removing information from semileptonic $b \rightarrow u\ell\nu$ decays [17]). Note that $|V_{ub}|_{\text{fit}}$ is quite close to the determination from exclusive decays and that $\text{BR}(B \rightarrow \tau\nu)_{\text{fit}}$ is *smaller* than the corresponding world average $((1.43 \pm 0.37) \times 10^{-4}$ [16]). The result in Eq. (10) is driven by the updated value for the B decay constant. The relatively low p-value [23] ($p = 15\%$) of the overall fit has been interpreted in terms of new physics in either K or in B_d mixing [6, 7, 8, 9, 10, 11].

Adopting the model independent parametrizations

$$|\varepsilon_K^{\text{NP}}| = C_\varepsilon |\varepsilon_K^{\text{SM}}|, \quad (12)$$

$$M_{12}^{d,\text{NP}} = e^{i\theta_d} M_{12}^{d,\text{SM}}, \quad (13)$$

$$\text{BR}(B \rightarrow \tau\nu)^{\text{NP}} = r_H \text{BR}(B \rightarrow \tau\nu)^{\text{SM}}, \quad (14)$$

where M_{12}^d is the matrix element of the effective Hamiltonian between B_d and $\bar{B}_d$ meson states and in the SM we have $(C_\varepsilon, r_H) = 1$ and $\theta_d = 0$. We obtain

$$C_\varepsilon = 1.28 \pm 0.14 \Rightarrow (2.0\sigma, p = 58\%), \quad (15)$$

$$\theta_d = -(4.0 \pm 1.8)^\circ \Rightarrow (2.2\sigma, p = 61\%), \quad (16)$$

$$r_H = 1.7 \pm 0.5 \Rightarrow (1.4\sigma, p = 29\%). \quad (17)$$

The above results are mutually exclusive in the sense that they are obtained by allowing either new physics in K , or in B_d mixing or in $B_u \rightarrow \tau\nu$ and point to a $\sim 2\sigma$ hint for new physics. The quoted p-values have to be compared with SM result ($p_{\text{SM}} = 15\%$). The lower p-value for new physics in $B \rightarrow \tau\nu$ indicates that the tension in the fit is only partially lifted by new contributions to $B \rightarrow \tau\nu$.

Removing semileptonic decays Recall that inclusive and exclusive $b \rightarrow (c, u)\ell\nu$ decays are tree-level Standard Model processes and, therefore most likely, are quite insensitive to the presence of new physics. The $|V_{cb}|$ constraint translates into a determination of the parameter A of the CKM matrix. Knowledge of the latter is critical in order to extract information from $|V_{ub}| \propto A$,




FIG. 2: *Upper panel:* interplay of the ε_K , ΔM_{B_s} and $\text{BR}(B \rightarrow \tau\nu)$ constraints. *Lower panel:* Unitarity triangle fit without semileptonic decays. The contour is obtained using ε_K , $B \rightarrow \tau\nu$, γ , ΔM_{B_s} and ΔM_{B_d} .

$\text{BR}(B \rightarrow \tau\nu) \propto A^2$ and $\varepsilon_K \propto A^4$ (see Eqs. (4,5)). The (ρ, η) regions allowed by each of these three observables is obtained *with* the inclusion of $|V_{cb}|$. Without any information on A these bands would cover the whole (ρ, η) plane. The main role of the determination of $|V_{ub}|$ is to limit the amount of new physics contributions to the phase of B_d mixing; in fact, an upper limit on $|V_{ub}|$ implies an upper limit on $S_{\psi K} = \sin 2(\beta + \theta_d)$ [24, 25]. Presently inclusive and exclusive determinations of $|V_{cb}|$ and $|V_{ub}|$ differ at the 2σ level (see Table I). The exclusion of the $|V_{ub}|$ constraint is not critical any longer to the presence of the 2σ effects in Eqs. (15,16) as emphasized recently in [7]. In particular the prediction that we obtain for the B_d mixing phase in the no- V_{ub} scenario reads $[\sin 2\beta]_{\text{fit}} = 0.840 \pm 0.056$ deviating by 2.8σ from its direct determination. On the other hand, $|V_{cb}|$ appears to be central: employing only its exclusive (inclusive) determination, the 2.0σ significance of the extraction of C_ε shifts to 2.5σ (1.6σ); similarly the 2.2σ effect in B_d mixing shifts to 2.9σ (1.4σ).

We now come to elaborating on the new approach that we are advocating here in which no use of semi-leptonic decays will be made. Note that the critical issue is the determination of A from $|V_{cb}|$. We find that the interplay of ε_K , $\text{BR}(B \rightarrow \tau\nu)$ and ΔM_{B_s} results in a fairly strong constraint on the (ρ, η) plane even without using semileptonic decays at all. A simple way to understand




FIG. 3: 95% C.L. bounds in the $(\tan\beta, m_{H^+})$ plane. The shaded region is excluded in the 2HDM. The dotted (dashed) line shows how this region shifts in two MSSM scenarios with $\epsilon_0 = -0.01$ (0.01) (the arrows indicate the excluded region).

this result is to use Eqs. (1-5) to eliminate A and write:

$$|\varepsilon_K| \propto \widehat{B}_K (f_{B_s} \widehat{B}_s^{1/2})^{-4} \quad (18)$$

$$|\varepsilon_K| \propto \widehat{B}_K \text{BR}(B \rightarrow \tau\nu)^2 f_B^{-4} \quad (19)$$

where for simplicity we kept only the dominant contributions to ε_K (proportional to A^4) and did not explicitly write the dependence of ε_K on ρ , η and all other quantities that are irrelevant to the error budget. Eqs. (18) and (19) show that the $|V_{cb}|$ constraint can be effectively replaced by either $f_{B_s} \widehat{B}_s^{1/2}$ or $\text{BR}(B \rightarrow \tau\nu) \times f_B^{-2}$. In the upper and lower panels of Fig. 2 we show respectively the anatomy of the ε_K , ΔM_{B_s} and $\text{BR}(B \rightarrow \tau\nu)$ constraints and the complete fit of the unitarity triangle in absence of semileptonic decays. The fit results for $|V_{ub}|$ and $\text{BR}(B \rightarrow \tau\nu)$ do not deviate significantly from Eqs. (9-10); the extracted value of $[\sin 2\beta]_{\text{fit}} = 0.811 \pm 0.074$ deviates by 1.8σ from its direct determination. It is interesting to observe that the result $|V_{cb}|_{\text{fit}} = (43.2 \pm 0.9) \times 10^{-3}$ is slightly larger than the average we quote in Table I: this is yet another manifestation of the tension between K and B_d mixing that we are observing in the fit. Finally, we note that $f_{B_s} \widehat{B}_s^{1/2}$ and $\xi = f_{B_s} \widehat{B}_s^{1/2} / f_{B_d} \widehat{B}_d^{1/2}$ are largely independent because they are affected by different lattice systematics and we average results from different lattice collaborations thereby reducing the possible correlation between statistical errors. A surprising outcome is the slight preference of the fit for new physics in B_d mixing. This can be seen by extracting C_ε , θ_d and r_H

$$C_\varepsilon^{\text{no}V_{qb}} = 1.23 \pm 0.30 \Rightarrow (0.8\sigma, p = 39\%), \quad (20)$$

$$\theta_d^{\text{no}V_{qb}} = -(8.4 \pm 4.6)^\circ \Rightarrow (1.8\sigma, p = 87\%), \quad (21)$$

$$r_H^{\text{no}V_{qb}} = 1.7 \pm 0.5 \Rightarrow (1.4\sigma, p = 64\%). \quad (22)$$

and noting that new physics in B_d mixing yields larger p -value than new physics in K mixing or in $B \rightarrow \tau\nu$.

As an illustration of the implications of these constraints we consider the impact on two Higgs dou-




FIG. 4: Reducing the errors on $f_{B_s} \widehat{B}_s^{1/2}$ and $\text{BR}(B \rightarrow \tau\nu)$.

blet models. Within these models the $r_H \neq 1$ result can be translated into a constraint on the mass of charged Higgs. In the type-II two Higgs Doublet Model (2HDM) and in the Minimal Supersymmetric SM (MSSM) we can write [26] $r_H = (1 - X_H)^2$ where $X_H = (\tan\beta m_{B^+}/m_{H^+})^2 / (1 + \epsilon_0 \tan\beta)$, $\tan\beta$ is the ratio of the vacuum expectation values of the Higgses that couple to up and down quarks and ϵ_0 summarizes supersymmetric corrections to the $\bar{u}bW^+$ vertex. In the 2HDM we have $\epsilon_0 = 0$; in the MSSM ϵ_0 does not vanish and typical values range at the 10^{-2} level. A full supersymmetric analysis of Eq. (22) is beyond the scope of this letter. In Fig. 3 we present the regions of the $(\tan\beta, m_{H^+})$ that are allowed at 95% C.L. for various values of ϵ_0 . In addition to the bounds implied by Eq. (22) we include also constraints from $B \rightarrow D\tau\nu$ [27] and $B \rightarrow X_s\gamma$ [28]. From the observation that X_H is always positive follows that the charged Higgs exchange can only reduce the $B \rightarrow \tau\nu$ branching ratio unless $X_H > 2$ implying a sign switch in the $B \rightarrow \tau\nu$ amplitude. Eq. (22) implies $X_H = (2.3 \pm 0.2) \vee (-0.3 \pm 0.2)$. At the 1σ level only the $X_H \sim 2$ solution is permitted (remember that $X_H > 0$) and the resulting allowed narrow band at low M_{H^+} or large $\tan\beta$, is, in turn, excluded by $B \rightarrow D\tau\nu$ data both in the 2HDM and in the MSSM (we follow the numerical analysis of Ref. [27]). At 95% C.L. the solution $X_H = 0$ opens up, corresponding to large M_{H^+} . In the 2HDM the $B \rightarrow X_s\gamma$ constraint implies $m_{H^+} > 295$ GeV [28]. In the MSSM, chargino loops contributions to the $b \rightarrow s\gamma$ amplitude can compensate charged Higgs effects: the bound on the charged Higgs depends strongly on the chosen point in the supersymmetric parameter space [29].

Let us now discuss the dominant sources of uncertainties in this analysis. In the following table we list the most relevant inputs, their errors and their impact on ε_K (as it follows from Eqs. (4), (18) and (19)):

$X :$	$\widehat{B}_K$	$ V_{cb} $	$f_{B_s} \widehat{B}_s^{1/2}$	$\text{BR}(B \rightarrow \tau\nu)$	f_B
$\delta X :$	4%	2.5%	6.9%	26%	5%
$\delta \varepsilon_K :$	4%	10%	27.6%	52%	20%

First of all, note that the impact of $\widehat{B}_K$ on the error is

subdominant. The use of the semileptonic $b \rightarrow c$ constraint results in a $\sim 10\%$ determination of ε_K , roughly half of the uncertainty obtained by employing only ΔM_{B_s} (i.e.: $f_{B_s} \hat{B}_s^{1/2}$). A calculation of $f_{B_s} \hat{B}_s^{1/2}$ at the 2.5% level would reduce the overall uncertainty on ε_K to 10%; a calculation at the 1% level would impact ε_K at the same level as $\hat{B}_K$. At first sight, the impact of $B \rightarrow \tau\nu$ seems irrelevant. Fortunately the non-trivial dependence of $\text{BR}(B \rightarrow \tau\nu)$ on ρ and η implies a certain degree of orthogonality between the constraints (18) and (19), as can be seen explicitly in the upper panel of Fig. 2. A numerical estimate of the impact of this constraint can be obtained by removing it from the fit and recalculating the overall p-value: we obtain $p = 43\%$, meaning that no hint of new physics is observed. The experimental uncertainty on the $B \rightarrow \tau\nu$ branching ratio is therefore an important ingredient of this analysis. Once the latter reaches the 10% level, improvements on f_B will be relevant as well. We summarize this discussion in Fig. 4 and in the following table:

δ_τ	δ_s	p_{SM}	$\theta_d \pm \delta\theta_d$	p_{θ_d}	$\theta_d/\delta\theta_d$
*26%	*6.8%	32%	$-(8.4 \pm 4.6)^\circ$	87%	1.8σ
*26%	2.5%	3.3%	$-(9.6 \pm 3.5)^\circ$	85%	2.7σ
*26%	1%	0.1%	$-(10.0 \pm 3.0)^\circ$	84%	3.4σ
10%	*6.8%	2%	$-(8.7 \pm 2.8)^\circ$	87%	3.1σ
3%	*6.8%	0.08%	$-(8.7 \pm 2.3)^\circ$	87%	3.8σ
10%	2.5%	0.1%	$-(9.2 \pm 2.5)^\circ$	84%	3.7σ
10%	1%	0.004%	$-(9.6 \pm 2.2)^\circ$	83%	4.3σ
3%	2.5%	0.004%	$-(9.1 \pm 2.1)^\circ$	84%	4.4σ
3%	1%	0.00009%	$-(9.4 \pm 1.9)^\circ$	82%	5.0σ

where $\delta_\tau = \delta\text{BR}(B \rightarrow \tau\nu)$, $\delta_s = \delta(f_{B_s} \hat{B}_s^{1/2})$ and * denotes the current uncertainties. The values $\delta_\tau = (10, 3)\%$ correspond to a super- B factory result with $(5, 50)\text{ab}^{-1}$ [30]. In the table we show the p-value of the SM fit and the result of the analysis of the scenario with new physics in B_d mixing (we display the new physics phase θ_d , the p-value of the new physics fit and its significance). We do not show the scenarios with new physics in K mixing or $B \rightarrow \tau\nu$ because they yield very low confidence levels and are, therefore, disfavored. From the inspection of the table, we conclude that even modest improvements in $f_{B_s} \hat{B}_s^{1/2}$ and/or $\text{BR}(B \rightarrow \tau\nu)$ will help enormously in isolating the presence of new physics in the unitarity triangle fit. For comparison, reducing the total uncertainty on $|V_{cb}|$ to the 1% level, yields $p_{\text{SM}} = 2.8\%$ corresponding to 2.7σ effects in either θ_d or C_ϵ .

Conclusions The tension in the standard fit of the unitarity triangle can be interpreted as a hint for new physics. The $\sim 2\sigma$ discrepancies in the extraction of $|V_{cb}|$ and $|V_{ub}|$ between inclusive and exclusive semileptonic B decays tend to cast doubt on the reliability of this conclusion. We investigated the removal of these

constraints from the fit. In contrast with the generally accepted statement that information on $|V_{cb}|$ is required in order to make meaningful use of ε_K , we showed, for the first time that, the combination of ε_K , ΔM_{B_s} and $\text{BR}(B \rightarrow \tau\nu)$ provide quite a stringent constraint in the (ρ, η) plane.

After removing information from semileptonic decays, we find that the tension in the unitarity triangle fit survives and can be translated into a 1.8σ hint for new physics in B_d mixing. The preference of the fit for new physics contributions to B_d mixing is caused by the $B \rightarrow \tau\nu$ constraint. This branching ratio is proportional to $|V_{ub}|^2$ and, as it can be seen from Fig. 2, points to a large value of $|V_{ub}/V_{cb}|$. This, in turn, favors a scenario with new physics in the B_d mixing phase.

Even modest improvements on $\text{BR}(B \rightarrow \tau\nu)$ and/or $f_{B_s} \hat{B}_s^{1/2}$ may push this tension above the 3σ level; if errors on both constraints are reduced simultaneously, $\delta_\tau = (10, 3)\%$ and $\delta_s = (2.5, 1)\%$, the effect reaches $(4 - 5)\sigma$. Note that improvements on $B \rightarrow \tau\nu$ require a super- B factory [31, 32, 33] while the reduction of δ_s is a purely theoretical (i.e. lattice) issue. Note also that lattice calculations of matrix elements relevant for decay constants or for (K^0, B_d, B_s) oscillations and therefore for δ_s , do not require momentum injection unlike the calculation of the semileptonic form factors and to that extent are simpler. However, we stress again that we are not suggesting abandoning the traditional approach with use of semileptonic decays, but rather in addition making concerted efforts towards improved lattice determination of $f_{B_s} \hat{B}_s^{1/2}$ and also of the $\text{BR}(B \rightarrow \tau\nu)$. These should provide valuable redundancy in our quest for new physics through flavor studies even in the LHC era. Finally we would like to stress that the main focus of the present letter is to propose a new clean strategy to implement simultaneously K and B_d mixing constraints on the (ρ, η) -plane and that our projections on the reach of this method depend solely on improving δ_τ and δ_s and are quite insensitive to the rest of the inputs summarized in Table I, in particular, the assumed errors in lattice computations.

This research was supported in part by the U.S. DOE contract No.DE-AC02-98CH10886(BNL).

-
- [1] N. Cabibbo, Phys. Rev. Lett., **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys., **49** 652 (1973).
 - [2] E. Gamiz et al., Phys. Rev., **D73**, 114502 (2006).
 - [3] D. J. Antonio *et al.*, Phys. Rev. Lett. **100**, 032001 (2008).
 - [4] C. Aubin et al., arXiv:0905.3947.
 - [5] C. Kelly [RBC/UKQCD], talk at Lattice09, <http://rhcpc.pku.edu.cn/workshop/lattice09/index.xml>.
 - [6] E. Lunghi and A. Soni, JHEP **09**, 053 (2007).
 - [7] E. Lunghi and A. Soni, Phys. Lett. **B666** 162 (2008).

- [8] A. J. Buras and D. Guadagnoli, Phys. Rev. **D78**, 033005 (2008).
- [9] A. J. Buras and D. Guadagnoli, arXiv:0901.2056.
- [10] E. Lunghi and A. Soni, arXiv:0903.5059.
- [11] J. Laiho et al., arXiv:0910.2928.
- [12] A. Lenz and U. Nierste, JHEP **06**, 072 (2007).
- [13] T. Aaltonen et al., Phys. Rev. Lett. **100**, 161802 (2008).
- [14] V. M. Abazov et al., Phys. Rev. Lett. **101**, 241801 (2008).
- [15] M. Bona et al., arXiv:0803.0659.
- [16] Heavy Flavor Averaging Group, update presented at LP2009, <http://www.slac.stanford.edu/xorg/hfag>
- [17] In this letter, for clarity, only constraints coming from $b \rightarrow (c, u)\ell\nu$ with $\ell = e, \mu$ are called direct determinations of $|V_{cb}|$ and $|V_{ub}|$.
- [18] J. Charles et al., Eur. Phys. J. **C41**, 1 (2005).
- [19] M. Ciuchini et al., JHEP **07**, 013 (2001).
- [20] M. Bona et al., arXiv:0906.0953.
- [21] R. T. Evans et al., PoS **LATTICE2008**, 052 (2008).
- [22] E. Gamiz et al., Phys. Rev. D **80**, 014503 (2009).
- [23] We remind the reader that the p-value is defined as the probability to obtain a minimum chi-square larger than the observed one, assuming the SM is correct (e.g. $p = 15\%$ means that there is a 15% chance that the observed result is explained by fluctuations).
- [24] W. Altmannshofer et al., arXiv:0909.1333.
- [25] A. J. Buras, arXiv:0910.1032.
- [26] W. S. Hou, Phys. Rev. D **48**, 2342 (1993).
- [27] U. Nierste et al., Phys. Rev. D **78**, 015006 (2008).
- [28] M. Misiak et al., Phys. Rev. Lett. **98**, 022002 (2007).
- [29] G. Barenboim et al., JHEP **0804**, 079 (2008).
- [30] T. Iijima, talk at “Hints for new physics in flavor decays”, KEK, Tsukuba, Japan, <http://belle.kek.jp/hints09>.
- [31] A. G. Akeroyd et al., hep-ex/0406071.
- [32] M. Bona et al., arXiv:0709.0451.
- [33] T. E. Browder et al., arXiv:0802.3201.